

FOR IMMEDIATE RELEASE Feb. 11, 2003.

## **Obstetricians Use Dubious Method In Attempt to Discredit Homebirth—**

Motives Questioned by Parents, Midwives, and Public Health Researchers

For the last six months, the American College of Obstetricians and Gynecologists (ACOG) has widely publicized a new study from Washington State that, prior to review by the wider scientific community, made homebirth appear unsafe. New analyses in the latest issue of *Obstetrics and Gynecology* confirm suspicions that major flaws in the study likely skewed the results. Parents, researchers, and midwives' associations across North America are concerned that, in spite of warnings about the inaccuracies, a prestigious medical organization promoted and continues to misrepresent the results of a poor study.

Long opposed to homebirth, ACOG has been unable to identify solid evidence that supports their position. With increasing numbers of studies using appropriate methods and enough births to make comparisons, it has become evident that proper care of healthy low risk women produces similar outcomes, whether they choose to give birth in home or hospital.

The new study, "Outcomes of Planned Home Birth In Washington State" (Pang, J. et al, *Obstet Gynecol.* 2002 August; 100(2): 253-9), was intended to compare neonatal outcomes for "planned" home births with those of planned hospital births. However, the study depends on birth certificate data that did not include information on the *intended* place of birth. The authors admitted that an unknown number of unplanned and unattended home births might have been included. Thus there were questions from the beginning about how reliable the study would be.

In the issue of *Obstetrics and Gynecology*, January 2003, letters to the editor compare the Washington study to other studies that used more sophisticated methods. These studies show that typically 8% to 10% of women who have planned home births attended by midwives move to the hospital during their labor. These transports are rarely considered emergencies. Statistics demonstrate they usually occur because of interest in speeding up the delivery with pain or hormonal medication and occasionally because of signs of potential fetal distress.

In contrast to the other studies, the Washington study reported only a 4.5% transport rate, much lower than expected with planned homebirth. This strongly indicates that the so-called "planned home births" included a large number of *unplanned* and *unattended* home births. These births are statistically more prone to bad outcomes, since there is no one trained to perform necessary procedures or recognize other care is needed in a timely fashion.

Although given opportunity, the investigators in Washington failed in their "Reply By The Authors" to respond to criticism about their abnormally low rate of transfer or to concerns about their misrepresentation of the homebirth literature. While the Washington study's conclusions are not likely valid even for Washington State, they definitely cannot be generalized beyond that state's borders.

Since the 1980s, researchers in the field have known the importance of distinguishing home births that are “planned” with a skilled attendant from those that occur at home by accident, without basic knowledge or equipment. In the Washington study, babies born in taxis, prematurely, or to mothers unaware or unable to access needed help could be included in the study as “planned” homebirths.

Licensed midwives in Washington State had brought flaws in the study design to the attention of the investigators prior to the initial presentation of the results. Among other indiscretions, they pointed out that a physician often signs the birth certificate when a baby who was born at home, without an attendant, is subsequently transferred to hospital. Thus the birth certificate data gives the false impression that the physician attended the home birth.

The midwives’ appeal to accurately represent the information was ignored. The press was given the story before publication in a medical journal, a protocol not normally followed in science circles. As a result, misleading headlines reached the public before the study’s methodology could be evaluated.

In contrast to the ACOG position, the American Public Health Association passed a resolution in 2001 to increase access to out-of-hospital birth attended by credentialed direct entry midwives. They based their decision on the weight of evidence about home birth demonstrated in the home birth studies carried out with better methodologies than were used in the Washington study.

**Press Release Issued By:**

Midwives Alliance of North America: [www.mana.org](http://www.mana.org)

The Canadian Association of Midwives: [www.canadianmidwives.org](http://www.canadianmidwives.org)

Citizens for Midwifery: [www.cfmidwifery.org](http://www.cfmidwifery.org)

North American Registry of Midwives: [www.narm.org](http://www.narm.org)

International Cesarean Awareness Network: [www.ican-online.org](http://www.ican-online.org)

Midwives and Mothers Watching Globally/Sages-Femmes et Mères Veillant Sur l’Humanité

The Association of Texas Midwives: [www.texasmidwives.com](http://www.texasmidwives.com)

The Global Birth Institute and Mother Nurture

American College of Domiciliary Midwives: [www.goodnewsnet.org](http://www.goodnewsnet.org)

**Contacts:**

1. Susan Hodges, President, Citizens for Midwifery [info@cfmidwifery.org](mailto:info@cfmidwifery.org) (888) 236-4880

2. Diane Holzer, President, Midwives Alliance of North America [Dhmidwife@compuserve.com](mailto:Dhmidwife@compuserve.com) (415) 721-7693

3. Kim Campbell, The Canadian Association of Midwives [kimcrm@shaw.ca](mailto:kimcrm@shaw.ca) (604) 859-0777.

4. Betty-Anne Daviss and Ken Johnson (researcher and epidemiologist) [midwife@istar.ca](mailto:midwife@istar.ca) and [Ken\\_LCDC\\_Johnson@hc-sc.gc.ca](mailto:Ken_LCDC_Johnson@hc-sc.gc.ca), h(613) 730-0282 or (613) 730-2323 Ottawa, Canada

## “Planned” Home Birth Study\* Misrepresented in the Press

The American College of Obstetricians and Gynecologists (ACOG) has publicized the Pang study\* with a press release that **misrepresents the study, misleads the public** about the actual conclusions and **omits mention of the study’s serious flaws**.

The authors themselves concluded that their study **should not be used to advise consumers** about the safety of planned home birth: “More light needs to be shed on this controversial topic before practitioners and expectant parents can be fairly counseled about the safety of planned home births.”

- The Pang study was based on birth registry data (birth certificates). Because birth certificate data does not include any information about the **intended site of birth**, the authors made **inappropriate assumptions** about which births were “planned” home births.
  - **No member of the research team had direct knowledge or experience with home birth**, and **they did not consult** with a single home birth expert or practitioner, so the authors **failed to consider** fundamental differences between home and hospital.
  - The authors admit that unplanned and unattended **births were wrongfully included** as “planned home births.”
  - The authors attributed to “planned home birth” perinatal deaths unrelated to the site of birth, and **failed to demonstrate any causal relationship between neonatal mortality and intended birth site**.
  - The authors **neglected to explain why their results contradict** every other study that has examined the safety of planned home births, a basic requirement of scientific studies.
- *The Pang study does not change the well-documented fact: **For a healthy low risk woman, a planned home birth attended by an experienced midwife (or physician) is as safe or safer than a doctor-attended birth in a hospital.***

\*Pang, JWY, et al “Outcomes of Planned Home Birth In Washington State,” *Obstetrics & Gynecology*, 2002 Aug;100(2):253-259

See other side for quotes from this study.

## **Can We Trust the Pang Study's Results?**

### **Here is what the authors say about their own study\*:**

"Because Washington State birth certificates do not identify which home births are planned, we defined planned home births ...." (Materials and Methods)

"This study has several limitations that are related to the reliance on birth certificate data. These include the potential for misclassifying unplanned home births ... and for misclassifying various outcomes and co-variates. In addition, data were missing for some potential confounders and effect modifiers." (Discussion)

"...misclassification of any unplanned home births as planned home births ... would result in inflated risk estimates of neonatal mortality and other outcomes for planned home births."

(Discussion)

"Several of the outcomes in this study may have been misclassified, namely respiratory distress requiring assisted ventilation for more than 30 minutes, prolonged labor, and postpartum hemorrhage." (Discussion)

"The likelihood of misclassification might be greater in a home setting than in a hospital ... the magnitude and direction of any such bias cannot be predicted and so caution should be used when interpreting the results for these outcomes." (Discussion)

"Nonetheless, more light needs to be shed on this controversial topic before practitioners and expectant parents can be fairly counseled about the safety of planned home births." (Discussion)

\*Pang, JWY, et al "Outcomes of Planned Home Birth In Washington State," *Obstetrics & Gynecology*, 2002 Aug;100(2):253-259

A basic literature review of previous studies,  
including one conducted in Washington State, demonstrates that

**HOME BIRTH IS SAFE FOR MOTHERS AND BABIES.**

Bibliography available upon request.

## Bibliography of Home Birth Studies

This bibliography focuses on industrialized countries only, primarily the United States and Great Britain. Citations are listed in reverse chronological order by publication date within the geographical settings. A complete list of citations, including abstracts, is available at <http://www.cfmidwifery.org>. Citations and abstracts were obtained from National Library of Medicine's Medline database.

### Overview:

Olsen O. "Meta-analysis of the safety of home birth." *Birth*, 1997 Mar;24(1):4-13; discussion 14-6. Department of Social Medicine, University of Copenhagen, Denmark.

### United States:

- Vedam, S. "Homebirth v. Hospital Birth: Questioning the Quality of Evidence for Safety." *Birth*, Mar 2003; 30:1:57-63. Department of Nurse Midwifery, Yale University, New Haven, CT, USA.
- Pang, JW, Heffelfinger JD, Huang GJ, Benedetti TJ, Weiss NS. "Outcomes of planned home births in Washington State: 1989-1996." *Obstet Gynecol* 2002 Aug;100(2):253-9. Comment in: *Obstet Gynecol* 2003 Jan;101(1):198; author reply 199-200. Department of Epidemiology, University of Washington School of Public Health and Community Medicine, Seattle, Washington 98195, USA. [jwpang@u.washington.edu](mailto:jwpang@u.washington.edu)
- DeVries R. "Midwifery in The Netherlands: vestige or vanguard?" *Med Anthropol* 2001;20(4):277-311. Department of Sociology and Anthropology, St Olaf College, Northfield, MN 55057, USA. [devries@stolaf.edu](mailto:devries@stolaf.edu)
- Murphy PA, Fullerton JT. "Measuring outcomes of midwifery care: development of an instrument to assess optimality." *J Midwifery Womens Health* 2001 Sep-Oct;46(5):274-84 Department of Obstetrics and Gynecology at Columbia University, NY, USA.
- Anderson RE, Anderson DA. "The cost-effectiveness of home birth." *J Nurse Midwifery* 1999 Jan-Feb;44(1):30-5 Dept. of Economics, Centre College, Danville, KY 40422, USA.
- Zeidenstein L. "Outcomes of intended home births in nurse-midwifery practice: a prospective descriptive study." *Journal of Nurse Midwifery*, 1999 Jan-Feb;44(1):80-1. Elizabeth Seton Childbearing Center, New York, New York, USA.
- Murphy PA. "Design and methodology in a community, practice-based research network: a study of nurse-midwifery home-birth practice." *Midwifery*, 1998 Sep;14(3):155-61. Columbia University Department of Obstetrics and Gynecology, New York, NY 10032, USA.
- Murphy PA, Fullerton J. "Outcomes of intended home births in nurse-midwifery practice: a prospective descriptive study." *Obstetrics and Gynecology*, 1998 Sep;92(3):461-70. Department of Obstetrics and Gynecology, Columbia University College of Physicians and Surgeons, New York, New York 10032, USA. [pam15@columbia.edu](mailto:pam15@columbia.edu)
- Clarke SC, Martin JA, Taffel SM. "Trends and characteristics of births attended by midwives." *Statistics Bulletin / Metropolitan Insurance Companies*, 1997 Jan-Mar;78(1):9-18.
- Anderson RE, Murphy PA. "Outcomes of 11,788 planned home births attended by certified nurse-midwives. A retrospective descriptive study." *Journal of Nurse Midwifery*, 1995 Nov-Dec;40(6):483-92.
- Declercq ER, Paine LL, Winter MR. "Home birth in the United States, 1989-1992. A longitudinal descriptive report of national birth certificate data." *Journal of Nurse Midwifery*, 1995 Nov-Dec;40(6):474-82. Boston University School of Public Health, Maternal and Child Health Program, MA 02118, USA.
- Janssen PA, Holt VL, Myers SJ. "Licensed midwife-attended, out-of-hospital births in Washington state: are they safe?" *Birth*, 1994 Sep;21(3):141-8.
- Sakala C. "Midwifery care and out-of-hospital birth settings: how do they reduce unnecessary cesarean section births?" *Social Science and Medicine*, 1993 Nov;37(10):1233-50. Health Policy Institute, Boston University, MA 02215.
- Declercq ER. "Where babies are born and who attends their births: findings from the revised 1989 United States Standard Certificate of Live Birth." *Obstetrics and Gynecology*, 1993 Jun;81(6):997-1004. Merrimack College, North Andover, Massachusetts.
- Duran AM. "The safety of home birth: the farm study." *American Journal of Public Health*, 1992 Mar;82(3):450-3. Department of Health, Commonwealth of the Northern Marianas Islands, Rota.

- Albers LL, Katz VL. "Birth setting for low-risk pregnancies. An analysis of the current literature." *Journal of Nurse Midwifery*, 1991 Jul-Aug;36(4):215-20. University of Medicine and Dentistry of New Jersey.
- Anderson R, Greener D. "A descriptive analysis of home births attended by CNMs in two nurse-midwifery services." *Journal of Nurse Midwifery*, 1991 Mar-Apr;36(2):95-103.
- Acheson LS, Harris SE, Zyzanski SJ. "Patient selection and outcomes for out-of-hospital births in one family practice." *Journal of Family Practice*, 1990 Aug;31(2):128-36. Department of Family Medicine, Case Western Reserve University, Cleveland, Ohio 44106.
- Morse JM, Park C. "Home birth and hospital deliveries: a comparison of the perceived painfulness of Parturition." *Research in Nursing and Health*, 1988 Jun;11(3):175-81. Faculty of Nursing, University of Alberta, Edmonton, Canada.
- Hinds MW, Bergeisen GH, Allen DT. "Neonatal outcome in planned v unplanned out-of-hospital births in Kentucky." *JAMA*, 1985 Mar 15;253(11):1578-82.
- Koehler NU, Solomon DA, Murphy M. "Outcomes of a rural Sonoma County home birth practice: 1976-1982." *Birth*, 1984 Fall;11(3):165-70.
- Declercq ER. "Out-of-hospital births, U.S., 1978: birth weight and Apgar scores as measures of outcome." *Public Health Reports*, 1984 Jan-Feb;99(1):63-73.
- Sullivan DA, Beeman R. "Four years' experience with home birth by licensed midwives in Arizona." *American Journal of Public Health*, 1983 Jun;73(6):641-5.
- Simmons R, Bernstein S. "Out-of-hospital births in Michigan, 1972-79: trends and implications for the safety of planned home deliveries." *Public Health Reports*, 1983 Mar-Apr;98(2):161-70.
- McClain C. "Women's choice of home or hospital birth." *Journal of Family Practice*, 1981 Jun;12(6):1033-8.
- Burnett CA 3d, Jones JA, Rooks J, Chen CH, Tyler CW Jr, Miller CA. "Home delivery and neonatal mortality in North Carolina." *JAMA*, 1980 Dec 19;244(24):2741-5.
- Dingley EF. "Birthplace and attendants: Oregon's alternative experience, 1977." *Women and Health*, 1979 Fall;4(3):239-53.
- Cameron J, Chase ES, O'Neal S. "Home birth in Salt Lake County, Utah." *American Journal of Public Health*, 1979 Jul;69(7):716-7.
- Anderson S, Bauwens E, Warner E. "The choice of home birth in a metropolitan county in Arizona." *JOGN Nursing*, 1978 Mar-Apr;7(2):41-6.
- Mehl LE. "Delivery in the home." *Comprehensive Therapy*, 1978 Mar;4(3):18-26.
- Mehl LE, Peterson GH, Whitt M, Hawes WE. "Outcomes of elective home births: a series of 1,146 cases." *Journal of Reproductive Medicine*, 1977 Nov;19(5):281-90.
- Long PJ, Jefferis C. "Home deliveries." *Frontier Nursing Services Quarterly Bulletin*, 1975 Spring;50(4):33-6.
- Lee FE, Glasser JH. "Role of lay midwifery in maternity care in a large metropolitan area." *Public Health Reports*, 1974 Nov-Dec;89(6):537-44.

#### **Great Britain:**

- Benjamin Y, Walsh D, Taub N. "A comparison of partnership caseload midwifery care with conventional team midwifery care: labour and birth outcomes." *Midwifery* 2001 Sep;17(3):234-40. Neonatal Unit, Leicester General Hospital, Gwendolyn Rd, Leicester.
- Longworth L, Ratcliffe J, Boulton M. "Investigating women's preferences for intrapartum care: home versus hospital births." *Health Soc Care Community* 2001 Nov;9(6):404-13. Health Economics Research Group, Brunel University, Oxbridge, UK.
- Chamberlain G, Wraight A, Crowley P. "Birth at home." *Practicing Midwife*, 1999 Jul-Aug;2(7):35-9. Obstetrics at Singleton Hospital, Swansea.
- Wraight A. "Home births. A report of the 1994 Confidential Enquiry by the National Birthday Trust." *Midwives* 1997 May;110(1312):104-6.
- Northern Region Perinatal Mortality Survey Coordinating Group. "Collaborative survey of perinatal loss in planned and unplanned home births." *BMJ: British Medical Association*, 1996 Nov 23;313(7068):1306-9.
- Davies J, Hey E, Reid W, Young G. "Prospective regional study of planned home births. Home Birth Study Steering Group." *BMJ: British Medical Association*, 1996 Nov 23;313(7068):1302-6. Maternity Survey Office, Newcastle upon Tyne.
- Arya R, Pethen T, Johanson RB, Spencer SA. "Outcome in low risk pregnancies." *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 1996 Sep;75(2):F97-102. Department of Paediatrics, North Staffordshire Hospital and University of Keele Stoke on Trent.
- Young G, Drife J. "Home or hospital birth?" *Practitioner*, 1992 Jun; 236(1515):672-4. National Perinatal Epidemiology Unit, Oxford.

- Ford C, Iliffe S, Franklin O. "Outcome of planned home births in an inner city practice." *BMJ: British Medical Association*, 1991 Dec 14;303(6816):1517-9. Department of Primary Health Care, Whittington Hospital, London.
- Tew, M. "Place of birth and perinatal mortality." *Journal of the Royal College of General Practitioners*, 1985 Aug;35(277):390-4.
- Campbell R, Davies IM, Macfarlane A, Beral V. "Home births in England and Wales, 1979: perinatal mortality according to intended place of delivery." *British Medical Journal (Clin Res Ed)*, 1984 Sep 22;289(6447):721-4.
- Shepperdson B. "Home or hospital birth? A study of women's attitudes." *Health Visitor*, 1983 Nov;56(11):405-6.
- Davis JA. "The place of birth." *Archives of Disease in Childhood*, 1982 Jun;57(6):406-9.
- Tew M. "Perinatal mortality: is home a safer place?" *Health and Social Service Journal*, 1980 May 30;90(4695):702-5.
- Barry CN. "Home versus hospital confinement." *Journal of the Royal College of General Practitioners*, 1980 Feb;30(211):102-7.
- Tew M. "The safest place of birth: further evidence." *Lancet*, 1979 Jun 30;1(8131):1388-90.
- Coates M. "Safety of domiciliary midwifery." *Lancet*, 1968 Nov 30;2(7579):1193.
- Law RG. "Domiciliary confinement in three high risk obstetric groups." *Proceedings of the Royal Society of Medicine*, 1968 Oct;61(10):1029-32.

**Other Countries: Includes The Netherlands, Canada, and Switzerland**

- Janssen PA, Lee SK, Ryan EM, Etches DJ, Farquharson DF, Peacock D, Klein MC. "Outcomes of planned home births versus planned hospital births after regulation of midwifery in British Columbia." *CMAJ* 2002 Feb 5;166(3):315-23. Comment in: *CMAJ*. 2002 Feb 5;166(3):335-6. Centre for Community Health and Health Evaluation Research, BC Research Institute for Children's and Women's Health, Vancouver.  
[pjanssen@interchange.ubc.ca](mailto:pjanssen@interchange.ubc.ca)
- Viisainen K. "Negotiating control and meaning: home birth as a self-constructed choice in Finland" *Soc Sci Med* 2001 Apr;52(7):1109-21. University of Helsinki, Department of Public Health, Helsinki, Finland.  
[kirsi.viisainen@stakes.fi](mailto:kirsi.viisainen@stakes.fi)
- Wieggers TA, van der Zee J, Kerssens JJ, Keirse MJ. "Variation in home-birth rates between midwifery practices in The Netherlands." *Midwifery* 2000 Jun;16(2):96-104. NIVEL (Netherlands Institute of Primary Health Care), PO Box 1568, 3500 BN Utrecht, The Netherlands.
- Harris G. "Homebirth and independent midwifery." *Aust Coll Midwives Inc J* 2000 Jul;13(2):10-6
- Andersson SW, Niklasson A, Lapidus L, Hallberg L, Bengtsson C, Hulthen L. "Sociodemographic characteristics influencing birth outcome in Sweden, 1908-1930. Birth variables in the Population Study of Women in Gothenburg." *J Epidemiol Community Health* 2000 Apr;54(4):269-78. Department of Clinical Nutrition, Goteborg University, Sweden.
- De Reu PA, Nijhuis JG, Oosterbaan HP, Eskes TK. "Perinatal audit on avoidable mortality in a Dutch rural region: a Retrospective study." *Eur J Obstet Gynecol Reprod Biol* 2000 Jan;88(1):65-9. Verloskundig Centrum Midden Brabant, Boxtel Nijmegen University, The Netherlands.
- Morison S, Percival P, Hauck Y, McMurray A. "Birthing at home: the resolution of expectations." *Midwifery* 1999 Mar;15(1):32-9. Edith Cowan University, School of Nursing, Pearson, WA, Australia.
- Gulbransen G, Hilton J, McKay L, Cox A. "Home birth in New Zealand 1973-93: incidence and mortality." *N Z Med J* 1997 Mar 28;110(1040):87-9. Comment in: *N Z Med J*. 1999 Dec 10;112(1101):475. Auckland Home Birth Association.
- Wieggers TA, Keirse MJ, van der Zee J, Berghs GA. "Outcome of planned home and planned hospital births in low risk pregnancies: prospective study in midwifery practices in The Netherlands." *BMJ: British Medical Association*, 1996 Nov 23;313(7068):1309-13. NIVEL (Netherlands Institute of Primary Health Care), Utrecht, Netherlands.
- Ackermann-Liebrich U, Voegeli T, Gunter-Witt K, Kunz I, Zullig M, Schindler C, Maurer M. "Home versus hospital deliveries: follow up study of matched pairs for procedures and outcome. Zurich Study Team." *BMJ: British Medical Association*, 1996 Nov 23;313(7068):1313-8. Institute for Social and Preventive Medicine, University of Basle, Switzerland.
- Tyson H. "Outcomes of 1001 midwife-attended home births in Toronto, 1983-1988." *Birth* 1991 Mar;18(1):14-9.
- Abernathy TJ, Lentjes DM. "Planned and unplanned home births and hospital births in Calgary, Alberta, 1984-87." *Public Health Reports*, 1989 Jul-Aug;104(4):373-7. Calgary Health Services, Alberta.

## Obstetricians Use Dubious Method to Discredit Homebirth — How Do the Numbers Stack Up?

For the last six months, the American College of Obstetricians and Gynecologists (ACOG) has widely publicized a new study from Washington State that makes homebirth appear unsafe. Unfortunately, ACOG promoted “*Outcomes of Planned Home Birth In Washington State*” (by Pang et al, 2002) before it could be reviewed by the wider scientific community. A closer look reveals major flaws in the study skewed that the results.

The great majority of studies of healthy women who choose home-based birth services under the care of an experienced birth attendant document safe outcomes for both mothers and their babies. This newest study from Washington State was intended to compare neonatal outcomes for “planned” home births with those of planned hospital births. However, it used birth certificate data which did not include information on the *intended* place of birth. The authors admitted that an unknown number of unplanned and unattended home births may have been included. This and other factors make the study’s conclusions unreliable.

US total births per year: 4 million (Martin, Joyce, et al, “Births Final Data for 2001,” *National Vital Statistics Reports*, Vol 51, No. 5, December 18, 2002, p. 1)

Washington study total births: 16,726

National neonatal mortality: about 2 per 1,000 live births for home, hospital or birth centers (excluding congenital anomalies, including births ending in c-sections). (Rooks JP. Meta-analysis of the safety of home birth. *Birth* 1997;24:4-13)

Neonatal mortality in the Washington Study:

Total for home and hospital groups: 38

Total excluding congenital anomalies: 19

Home birth cohort: 12, or 1.95/1000 live births (including births ending in c-sections)

Hospital cohort: 7, or 0.7/1000 live births (NOT including births ending in c-sections)

National Cesarean /operative rate: 30%

WA study home birth cohort total Cesarean /operative rate: less than 4%

WA study hospital cohort total Cesarean /operative rate: none – these births were excluded from the study

Home to hospital transport rates reported in most home birth studies: 8 to 10%

Home to hospital transport rates reported in WA Study: about 4%, suggesting a substantial number of unplanned, unattended home births were included in the home birth cohort. (Johnson KC. Letter to the Editor. *Obs & Gyn.* 2003;101:1)

- The Washington Study neonatal mortality numbers are extremely small. Just one or two mistaken or misclassified cases could change the results.
- No chart reviews were conducted to clarify or verify the birth certificate data or determine any causal relationship between neonatal outcome and intended place of birth.
- The study design ignored the increased cost and maternal complications caused by the very high rate of medical interventions associated with “planned hospital birth”, such as Pitocin-accelerated labors, narcotics, episiotomies, forceps and 20 percent-plus Cesarean section rate.
- The hospital and home birth cohorts were not comparable since the results of intended hospital births that ended in Cesarean sections were not included, but intended home births that ended in Cesareans sections were included.
- Cesarean deliveries are riskier for mothers and babies and can result in maternal-infant mortality in subsequent pregnancies.